

# Gravitational Waves from Holographic Neutron Star Mergers

Christian Ecker



**Universiteit Utrecht**

Amsterdam – 25 October 2019

Based on arXiv:1908.03213

with Matti Järvinen, Govert Nijs and Wilke van der Schee

# Neutron star merger with holographic EoS

# Outline

1. Introduction
2. Holographic model
3. Merger simulations
4. Results
5. Summary and Outlook


# 1. Introduction

# Neutron stars

- ▶ Neutron stars (NSs) are born in supernova explosions of massive ( $8 - 25M_{\odot}$ ) main sequence stars.
- ▶ Densest astrophysical objects which are not black holes (BHs)  
 $M \approx 1.4M_{\odot}$ ,  $R \approx 10\text{km}$ ,  $\rho \approx 2 - 5\rho_0$   
(nuclear saturation mass-energy density  $\rho_0 = 2.5 \cdot 10^{14}\text{g cm}^{-3}$ )
- ▶ Can have huge magnetic fields (magnetars)  $B \approx 10^{15}\text{G}$   
(cf. earth:  $B_{\oplus} \approx 0.6\text{G}$ , RHIC:  $B_{HIC} \approx 10^{18}\text{G}$ )
- ▶ Some rotate extremely fast (pulsars)  $\nu_{\text{rot}} \lesssim 1\text{ms}^{-1}$   
first detection in 1967 as pulsar = rapidly rotating and highly magnetized NS  
record holder: PSR J1748-2446ad ( $v_R \approx 0.24c$ )
- ▶ Neutron stars are cool:  $T$  in keV range  
 $T \ll \Lambda_{QCD} \implies T = 0$  good approx., located at bottom of QCD phase diagram.  
neutrino cooling: from  $10^{11}\text{K} \approx 10\text{MeV}$  after supernova to keV within days.  
shock heating: finite temperature contributions become important during merger.

# Neutron star binaries

- ▶ Binary NS systems are most likely formed in main-sequence star binaries<sup>1</sup>, less likely by dynamical capture.
- ▶ First discovery in 1974: Hulse-Taylor binary pulsar (PSR B1913+16), successful test of Einstein gravity and indirect proof for gravitational waves (GWs).
- ▶ Known binary NS systems typically have:  
[J.M. Lattimer 2012, see also [www.stellarcollapse.org](http://www.stellarcollapse.org)]
  - ▶ Masses around  $M_{1,2} \approx 1.4M_{\odot}$
  - ▶ Mass ratio  $q \equiv M_1/M_2 \approx 1$
  - ▶ Several million years of inspiral phase, i.e. enough time to cool down by neutrino cooling and circularize orbits by GW emission.




[picture: Weisberg, Taylor  
[astro-ph/0407149](https://arxiv.org/abs/astro-ph/0407149)]

<sup>1</sup>Most stars (up to  $\approx 85\%$ ) are actually in binary systems. (see [www.atnf.csiro.au](http://www.atnf.csiro.au))

# Neutron star mergers

- ▶ Significant sources of gravitational radiation.
- ▶ Microscopic properties of dense matter encoded in GW and EM signal.
- ▶ Likely the origin of short gamma-ray bursts (SGRBs)
- ▶ and of heavy elements like gold, platinum, or uranium.





[picture: Baiotti, Rezzola arXiv:1607.03540]

(SMNS:  $M_{\text{TOV}} < M < M_{\text{max}} \approx 1.2M_{\text{TOV}}$ , HMNS:  $M > M_{\text{max}}$ )

# Neutron star equation of state

- ▶ Equation of state (EoS)  $p(\epsilon, T, \dots)$  required to close EOMs in fluid description of neutron stars.
- ▶ EoS of neutron stars is currently unknown because:
  - ▶ Nuclear matter models typically only reliable for  $\rho \lesssim \rho_0$ .
  - ▶ First principle QCD calculations at relevant densities not feasible:
    - ▶ pQCD only at asymptotically large densities and/or  $T$ .
    - ▶ Lattice QCD has sign problem at finite densities.





# Constraints on the EoS

From theory:

- By causality the speed of sound has to satisfy:  $c_s^2 = \left(\frac{\partial p}{\partial \epsilon}\right)_s < 1$ .
- QCD is asymptotically free at large densities:  $c_s^2 \rightarrow 1/3$ .

From observation:

- NS-white dwarf binary PSR J0348+0432:  $M_{\max} > 2.01 \pm 0.04 M_\odot$ .  
[Antoniadis et. al. arXiv:1304.6875]
- GW170817 constrains tidal deformability:  $\Lambda = \frac{2}{3} k_2 (c^2 R / (Gm))^5 \lesssim 600$ .  
[LIGO/Virgo: arXiv:1710.05832, arXiv:1805.11579, arXiv:1805.11581]



[Annala, Gorda, Kurkela, Nättila, Vuorinen arXiv:1903.09121, left plot by T. Gorda]

## 2. Holographic Model

# Holographic model: V-QCD

Bottom-up construction with dynamical quark and gluon sectors and potentials tuned to mimic QCD.

Two building blocks of Veneziano QCD (V-QCD):

1. Improved holographic QCD (IHQCD) for gluon sector (dilaton gravity).  
[Gürsoy, Kiritsis, Nitti; Gubser, Nellore]
2. Sen-like tachyonic Dirac-Born-Infeld (DBI) actions for flavor sector and chiral symmetry breaking via tachyon condensation.  
[Bergman, Seki, Sonnenschein; Dhar, Nag; Bigazzi, Casero, Cotrone, Kiritsis, Paredes; Gürsoy, Kiritsis, Nitti; Iatrakis, Kiritsis, Paredes]

Consider 1. + 2. in the Veneziano limit with full backreaction:

$$N_c \rightarrow \infty \text{ and } N_f \rightarrow \infty \text{ with } x \equiv N_f/N_c \text{ fixed}$$


[Järvinen, Kiritsis arXiv:1112.1261]

Add probe baryons: simple approximation with homogeneous bulk soliton. One free parameter:  $b$  = coupling between baryon and chiral condensate.

[Ishii, Järvinen, Nijs arXiv:1903.06169]

# Phase diagram

- ▶ Baryons appear at medium  $\mu$  in the confined phase
- ▶ Nontrivial nuclear and quark matter EoS from the same model
- ▶ Free parameter of the holographic model  $b$  chosen to set the correct  $\mu_c$  at vacuum-baryon transition



[Ishii, Järvinen, Nijs arXiv:1903.06169]

# Hybrid Equation of State (I)

Construction of the hybrid EoS:


- ▶ In the crust and for low densities (up to  $1.5 - 2\rho_0$ ) we use the SLy nuclear matter EoS. Here the nuclear matter EoS is expected to be more reliable than the holographic model.  
[Haensel, Pichon nucl-th/9310003, Douchin, Haensel astro-ph/0111092]
- ▶ For dense nuclear matter the baryonic V-QCD is matched<sup>2</sup> to SLy.
- ▶ Baryon onset in SLy part  $\implies$  freedom in holographic parameter  $b$ .
- ▶ Changing  $b$  shifts  $\mu_{\text{match}}$   $\implies$  more/less SLy at low densities.
- ▶ At high densities the V-QCD EoS transitions to the quark phase  
 $\implies$  same model for holographic baryon and quark phase.

---

<sup>2</sup>Matching point  $\mu_{\text{match}}$  and normalization  $c_b$  of the action are chosen to give continuous  $p(\mu)$  and  $p'(\mu)$  at matching point.

# Hybrid Equation of State (II)

- ▶ Second order phase transition at matching between SLy and VQCD.
- ▶ Strong<sup>3</sup> first order nuclear to quark matter phase transition.
- ▶ LIGO observation GW170817 constrains  $b \gtrsim 10.45 \rightarrow \Lambda_{1.4} \approx 680$ .




<sup>3</sup>The latent heat at the transition is sizable (for  $b = 10.5$ :  $\Delta\epsilon = 920 \text{ MeV}/\text{fm}^3$ ).

# Mass-radius relation

- Large values of  $b \gtrapprox 10.65$  are ruled out by  $2.01 M_{\odot}$ -bound from PSR J0348+0432 and PSR J1614-2230.

[Demorest et. al. arXiv:1010.5788, Antoniadis et. al. arXiv:1304.6875]

- Allowed values in the holographic model:  $10.45 \lesssim b \lesssim 10.65$ .



$b$	$R_{1.4}$ km	$M_{\max}$ $M_{\odot}$
10.45	13.0	2.25
10.50	12.6	2.17
10.55	12.3	2.10
10.60	12.1	2.04
10.65	12.0	2.00
SLy	11.7	2.05

### 3. Merger Simulations

# Simulating neutron star mergers (I)

Have to evolve the 4D general relativistic hydrodynamics equations:

$$R_{\mu\nu} - \frac{1}{2} R g_{\mu\nu} = 8\pi G_N T_{\mu\nu}, \quad \nabla_\mu T^{\mu\nu} = 0, \quad \nabla_\mu J^\mu = 0, \quad p = p(\epsilon),$$
$$ds^2 = -(\alpha^2 - \beta^i \beta_i) dt^2 + 2\beta_i dx^i dt + \gamma_{ij} dx^i dx^j,$$

with initial conditions (spacetime and fluid distribution) modelling a NS binary system:

- ▶ Use pseudo spectral code LORENE<sup>4</sup> to generate initial data.

[Gourgoulhon, Grandclement, Taniguchi, Marck, Bonazzola arXiv:gr-qc/0007028]

- ▶ Assumes helical symmetry = circularity of the BNS orbit, no GWs.
- ▶ Use non-spinning ICs for the fluid distribution.
- ▶ Initial distance  $D$  must be sufficiently large to justify circular orbit approximation, but not too large (reasonable inspiral time).  
Use  $D = 45\text{km}$ , gives between three ( $1.4M_\odot$ ) and six ( $1.3M_\odot$ ) orbits.

---

<sup>4</sup>Langage Objet pour la RElativite NumeriquE, see <http://www/lorene.obspm.fr>

# Simulating neutron star mergers (II)

Einstein equations need to be solved in strongly hyperbolic<sup>5</sup> form.

- ▶ Use Conformal and Covariant Z4 (CCZ4) formulation  
[Alic, Bona-Casas, Bona, Rezzolla, Palenzuela]
- ▶ Lapse  $\alpha$  and shift  $\beta^i$  fixed by singularity avoiding gauge conditions.
- ▶ Implemented in the Einstein Toolkit.  
[Löfler et. al., arXiv:1111.3344, <http://einstoolkit.org>]
- ▶ Hydro equations solved with WhiskyTHC<sup>6</sup>.  
[Radice, Rezzolla arXiv:1206.6502]

---

<sup>5</sup>The weakly hyperbolic ADM formulation (metric  $\gamma_{ij}$  and the extrinsic curvature  $K_{ij}$  as dynamical variables) does not lead to stable time evolution.

<sup>6</sup>see D. Radice's homepage: <https://www.astro.princeton.edu/~dradice/index.html>

# Spacetime discretization

- ▶ Grid must be large enough to extract GW in "asymptotic" region.
- ▶ Fine enough to resolve the NS, merger dynamics and BH formation.  
    ⇒ Adaptive mesh refinement (AMR) required.
- ▶ We use a cubic grid of  $\approx 3025\text{km}$  ( $\approx 70D$ ) side length.
- ▶ Reflection symmetry about  $xy$ -plane.
- ▶ Six "refinement levels":  
    Coarse mesh ( $\Delta h_0 \approx 23.5\text{km}$ ) in the asymptotic region.  
    Finest meshes ( $\Delta h_6 \approx 365\text{m}$ ) centered at the neutron stars.  
    ⇒  $\approx 5 \cdot 10^6$  gridpoints and  $\approx 5 \cdot 10^5$  time steps!

# Supercomputing

- ▶ To simulate neutron stars one needs supercomputing.
- ▶ Pilot project (500.000 CPUh) on Dutch supercomputer Cartesius.  
[<https://userinfo.surfsara.nl/systems/cartesius>]
- ▶ Typical simulation costs 10.000 CPUh on 100 CPUs and takes ≈ 4 days.



# Extracting Waveforms

- ▶ Extract Newman-Penrose scalar  $\psi_4$  from simulation

$$\psi_4 := C_{\mu\nu\rho\sigma} k^\mu \bar{m}^\nu k^\rho \bar{m}^\sigma, \quad \text{null tetrade: } \{l^\mu, k^\mu, m^\mu, \bar{m}^\mu\}.$$

- ▶ The GW polarization amplitudes ( $h_+, h_\times$ ) are related to  $\psi_4$  via  
[see e.g. book by M. Alcubierre (2005)]

$$\ddot{h}_+ - i\ddot{h}_\times = \psi_4 = \sum_{\ell=2}^{\infty} \sum_{m=-\ell}^{\ell} \psi_4^{\ell m} {}_{-2}Y_{\ell m}(\theta, \varphi).$$

- ▶ Consider only the dominant  $\ell = m = 2$  mode

$$h_{+,\times} = \sum_{\ell=2}^{\infty} \sum_{m=-\ell}^{\ell} h_{+,\times}^{\ell m} {}_{-2}Y_{\ell m}(\theta, \varphi) \approx h_{+,\times}^{22} {}_{-2}Y_{22}(\theta, \varphi),$$

and assume optimal orientation ( ${}_{-2}Y_{22}(0, 0) = \frac{1}{2}\sqrt{5/\pi}$ ) of the merger with respect to the detector.

- ▶ Extrapolate signal to detector, assume distance of 40Mpc, i.e. same luminosity distance as GW170817.

# Power Spectral Density


Post-merger power spectral density (PSD) has typical three peak structure.

$$\tilde{h}(f) \equiv \sqrt{\frac{|\tilde{h}_+(f)|^2 + |\tilde{h}_{\times}(f)|^2}{2}}, \quad \tilde{h}_{+,\times}(f) \equiv \int h_{+,\times}(t) e^{-i2\pi ft} dt.$$

Characteristic frequencies  $f_1$ ,  $f_2$ ,  $f_3$  contain information about EoS.

[Takami, Rezzolla, Baiotti arXiv:1403.5672]



Strategy: simulate predictions for  $f_1$ ,  $f_2$ ,  $f_3$  and compare to future observations.



## 4. Results



# High mass binary

- ▶ Equal mass binary  $M_1 = M_2 = 1.5M_\odot$  with  $b = 10.5$  SLy+VQCD EoS.
- ▶ Prompt collapse to BH with dilute matter torus  $\rightarrow$  trivial post-merger PSD.
- ▶ Densities are not high enough to create quark matter outside BH.





# Intermediate mass binary

- ▶ Equal mass binary  $M_1 = M_2 = 1.4M_{\odot}$  with  $b = 10.5$  SLy+VQCD EoS.
- ▶ Formation of a short lived ( $\approx 7.8\text{ms}$ ) HMNS  $\rightarrow$  non-trivial PSD.




# Low mass binary

- ▶ Equal mass binary  $M_1 = M_2 = 1.3M_{\odot}$  with  $b = 10.5$  SLy+VQCD EoS.
- ▶ Formation of a long lived ( $> 40\text{ms}$ ) HMNS  $\rightarrow$  non-trivial PSD.




# Mass dependence of the Power Spectral Density



$M[M_\odot]$	EoS	$b$	$f_1[\text{kHz}]$	$f_2[\text{kHz}]$	$f_3[\text{kHz}]$
1.30	SLyVQCD105	10.5	1.93	2.53	3.77
1.35	SLyVQCD105	10.5	1.95	2.60	3.53 (3.90)
1.40	SLyVQCD105	10.5	2.03	2.89	3.82
1.50	SLyVQCD105	10.5	—	—	—


# EoS dependence of the Power Spectral Density



$M[M_\odot]$	EoS	$b$	$f_1$ [kHz]	$f_2$ [kHz]	$f_3$ [kHz]
1.30	SLyVQCD105	10.5	1.93	2.53	3.77
1.30	SLyVQCD106	10.6	2.15	2.80	3.70 (4.06)
1.30	SLy	-	2.21	3.19	4.24

# Universality

- ▶ Frequency  $f_1$  as function of compactness  $\mathcal{C} = M/R$  shows universal behaviour, i.e. results for different EoS fall on one universal curve.
- ▶ V-QCD EoS for  $b = 10.5$  gives  $f_1$  close to universal curve,  $b = 10.6$  slightly off (?)  $\implies$  more analysis needed.



[plot from Takami, Rezzolla, Baiotti arXiv:1403.5672]

## 5. Summary and Outlook

# Summary

- ▶ Constructed “realistic” hybrid holographic EoS that satisfies all currently known theoretical and observational constraints.
- ▶ First NS merger simulations using input from holography.
- ▶ No (meta-)stable quark matter outside a BH horizon  
    ⇒ phase transition leads to prompt collapse.
- ▶ Predictions for characteristic peaks in power spectral densities  
    ⇒ new connection between holography and observation!
- ▶ Features in post merger power spectral densities (location of peaks) expected to be accessible by 3rd generation detectors.

# Outlook

- ▶ Increase resolution to improve accuracy of waveforms and PSD (costly!).
- ▶ Include finite  $T$  effects in EoS → temperature driven de-confinement phase transition.
- ▶ Spin, non-equal mass binaries, NS-BH merger, magnetic fields.
- ▶ Improve holographic model:
  - ▶ go beyond homogeneous baryon ansatz: numerical solutions for inhomogeneous solitons.
  - ▶ include magnetic field dependence in the EoS.
  - ▶ systematic study with different nuclear matter EoSs at low density.
- ▶ Only first step towards holographic GW model building, lots of possibilities and challenges!

## 6. Backup

# V-QCD without baryons (I)

Consider first the non-baryonic V-QCD action, whose solutions will serve as background for the probe baryons

$$S_{\text{V-QCD}}^{(0)} = S_{\text{glue}} + S_{\text{DBI}}^{(0)}.$$

The gluon part is given by the IHQCD (dilaton gravity) action

$$S_{\text{glue}} = N_c^2 M^3 \int d^5x \sqrt{-g} \left[ R - \frac{4}{3} \frac{(\partial \lambda)^2}{\lambda^2} + V_g(\lambda) \right],$$

where  $\lambda \equiv e^\phi \leftrightarrow \text{Tr} F^2$  ( $\approx g^2 N_c$  near the boundary) sources the 't Hooft coupling in YM theory, the dilaton potential is chosen<sup>7</sup> to mimic QCD

$$V_g(\lambda) = 12 \left[ 1 + V_1 \lambda + \frac{V_2 \lambda^2}{1 + \lambda/\lambda_0} + V_{\text{IR}} e^{-\lambda_0/\lambda} (\lambda/\lambda_0)^{4/3} \sqrt{\log(1 + \lambda/\lambda_0)} \right].$$

Finite  $T$  is implemented by homogeneous+isotropic black brane metric

$$ds^2 = e^{2A(r)} (-f(r) dt^2 + d\vec{x}^2 + f^{-1}(r) dr^2).$$

---

<sup>7</sup>E.g.  $V_1$  and  $V_2$  are fixed by requiring the UV RG flow of the 't Hooft coupling to be the same as in QCD up to two-loop order.

## V-QCD without baryons (II)

The flavor part is modelled by the tachyonic DBI-action<sup>8</sup>

$$S_{\text{DBI}}^{(0)} = -N_f N_c M^3 \int d^5x V_{f0}(\lambda) e^{-\tau^2} \sqrt{-\det [g_{ab} + \kappa(\lambda) \partial_a \tau \partial_b \tau + w(\lambda) F_{ab}]},$$
$$F_{rt} = \Phi'(r), \quad \Phi(0) = \mu,$$

where the tachyon  $\tau \leftrightarrow \bar{q}q$  controls chiral symmetry breaking.

Several potentials:  $\{V_g(\lambda), V_{f0}(\lambda), w(\lambda), \kappa(\lambda)\}$ , chosen to match pQCD in UV ( $\lambda \rightarrow 0$ ), qualitative agreement with QCD in IR ( $\lambda \rightarrow \infty$ ) and tuned to lattice QCD in the middle ( $\lambda \sim \mathcal{O}(1)$ ).

[For details see Appendix B of Ishii, Järvinen, Nijs arXiv:1903.06169]

Different solutions:

without/with horizon  $\leftrightarrow$  confined/deconfined phase

without/with tachyon  $\leftrightarrow$  chirally symmetric/chirally broken phase

---

<sup>8</sup>Without baryons we have a vectorial flavor singlet gauge field  $A^{(L/R)} = \mathbb{I}_f \Phi(r) dt$  giving nonzero charge density and chemical potential.

# Probe baryons in V-QCD

Each baryon maps to a solitonic “instanton” configuration of non-Abelian gauge fields in the bulk.

[Witten; Gross, Ooguri; ...]

Consider the full non-Abelian brane action  $S = S_{\text{DBI}} + S_{\text{CS}}$  where

[Bigazzi, Casero, Cotrone, Kiritis, Paredes; Casero, Kiritis, Paredes]

$$S_{\text{DBI}} = -\frac{1}{2} M^3 N_c \mathbb{T} r \int d^5 x V_{f0}(\lambda) e^{-\tau^2} \left( \sqrt{-\det \mathbf{A}^{(L)}} + \sqrt{-\det \mathbf{A}^{(R)}} \right),$$

$$\mathbf{A}_{MN}^{(L/R)} = g_{MN} + \delta_M^r \delta_N^r \kappa(\lambda) \tau'(r)^2 + \delta_{MN}^{rt} w(\lambda) \Phi'(r) + w(\lambda) \mathcal{F}_{MN}^{(L/R)}$$

gives the dynamics of the solitons.

The Chern-Simons term sources the baryon number for the solitons

$$S_{\text{CS}} = \frac{N_c}{8\pi^2} \int \Phi(r) e^{-b\tau^2} dt \wedge \left( F^{(L)} \wedge F^{(L)} - F^{(R)} \wedge F^{(R)} + \dots \right).$$

Non-Abelian DBI action only known to first few orders in  $\mathcal{F}^{(L/R)}$ : expand to second order on top of solution  $(g_{MN}, \Phi, \lambda, \tau)$  obtained from  $S_{V-QCD}^{(0)}$ .

# Homogeneous Baryon Ansatz

Set  $N_f = 2$  and consider the SU(2) Ansatz

[Rozali, Shieh, Van Raamsdonk, Wu]

$$A_L^i = -A_R^i = h(r)\sigma^i$$

Immediate consequence: baryon charge integrates to zero?

$$N_b \propto \int dr \frac{d}{dr} \left[ e^{-b\tau^2} h^3 (1 - 2b\tau^2) \right] \stackrel{?}{=} 0$$

However finite baryon number may can be realized by discontinuity of  $h$   
↔ smeared solitons in singular gauge.

[Ishii, Järvinen, Nijs, arXiv:1903.06169]

The free parameter  $b$  of the model is used to tune the baryon onset to its physical value in QCD.